

Sick Day Rules

It is important to note that Sick Day Rules require initiation from your Doctor, Nurse or Healthcare Practitioner.



When you are experiencing the following

- Vomiting or diarrhoea (unless only minor)
- Fevers, sweats and shaking (unless only minor)

It is easy to become dehydrated when you are ill. Dehydration may stress your body and may cause certain medications to cause problems

Pause taking the medicines your Healthcare Professional has ticked on the reverse of this card.

It is very important that you restart your medicine once you have recovered from your illness. This would normally be after 24 to 48 hours of eating and drinking normally. When you restart your medicine, just take them as normal. Do not take extra for the doses you have missed.

Please remember to keep yourself hydrated and try to weigh yourself daily.

Please contact your Heart Failure Team if:

- You find that you experience weight loss or gain of 1.5kg over two days
- Your blood pressure is unusually low for you
- You have been ill for more than 2 days
- You are in any doubt or have concerns

Medicines to Pause on Sick Days

- ACE Inhibitors & ARBs:** Medicine names ending in “pril” or ARB ends in “sartan”
- MRA:** e.g., Spironolactone, Eplerenone
- Diuretics:** e.g., Furosemide, Bumetanide, Torsemide, Bendroflumethiazide, Metolazone
- SGLT2 Inhibitors:** Medicine names ending in “gliflozin”
- ARNI:** e.g., Sacubitril Valsartan (Entresto)

Many thanks to

Professor Paul Kalra Consultant Cardiologist

Dr Rani Khatib Consultant Pharmacist in Cardiology

Carys Barton Heart Failure Nurse Consultant

For assisting with the authorship of this valuable patient resource.

Other Medications to Stop Taking

.....

.....

.....

.....



Help us make heart failure clearer for everyone

+44 (0) 1772 796542

hearts@pumpingmarvellous.org

www.pumpingmarvellous.org

LET'S
B.E.A.T
HEART FAILURE
TOGETHER